

Geophysical Research Letters[®]

RESEARCH LETTER

10.1029/2022GL102070

Rapid 21st Century Weakening of the Agulhas Current in a Warming Climate

Ruize Zhang^{1,2,3} , Shantong Sun⁴ , Zhaohui Chen^{1,2} , Haiyuan Yang^{1,2} , and Lixin Wu^{1,2}

¹Frontier Science Center for Deep Ocean Multispheres and Earth System (FDOMES), Physical Oceanography Laboratory, Ocean University of China, Qingdao, China, ²Laoshan Laboratory, Qingdao, China, ³Academy of the Future Ocean, Ocean University of China, Qingdao, China, ⁴Department of Earth, Ocean & Atmospheric Science, Florida State University, Tallahassee, FL, USA

Key Points:

- Climate models consistently project the Agulhas Current (AC) to weaken in the 21st century
- Changes in Indo-Pacific wind and Atlantic overturning both contribute to the projected AC weakening
- The AC becomes shallower in a warming climate mainly due to the weakening of the Atlantic overturning

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

S. Sun and Z. Chen,
ssun5@fsu.edu;
chenzhaohui@ouc.edu.cn

Citation:

Zhang, R., Sun, S., Chen, Z., Yang, H., & Wu, L. (2023). Rapid 21st century weakening of the Agulhas current in a warming climate. *Geophysical Research Letters*, 50, e2022GL102070. <https://doi.org/10.1029/2022GL102070>

Received 9 NOV 2022
 Accepted 11 FEB 2023

Abstract The Agulhas Current (AC) is a critical component of the global ocean circulation. Climate models consistently project the AC to decline in the 21st century. Previous studies typically attributed the weakening of AC to changes in Indian Ocean wind and/or a decline of the Indonesian Throughflow (ITF) transport. However, our analysis suggests that changes in local surface wind and ITF can only explain a portion of the model-simulated AC changes. Using a hierarchy of models, we show that the decline of the Atlantic Meridional Overturning Circulation (AMOC) strength in a warming climate could also contribute substantially to the AC weakening. Following a weakening of the AMOC, the baroclinic structure of the AC is also modified such that the AC is shoaled to a shallower depth, communicated between basins via Kelvin waves. Our results highlight the importance of remote processes in future western boundary current changes.

Plain Language Summary The Agulhas Current (AC) is one of the strongest currents in the ocean and plays an important role in the climate system. Climate models consistently project a rapid weakening of the AC during the 21st century. Traditionally, the weakening of the AC has been attributed to changes in local surface wind and the Indonesian Throughflow, which transports water from the Pacific Ocean to the Indian Ocean. However, our analysis suggests these changes cannot fully explain the AC changes in a warming climate. In this study, we show that remote processes in the North Atlantic could also contribute substantially to the AC weakening. In a warming climate, the Atlantic overturning circulation slows down and modifies the density structure of the Atlantic, which propagates into the Indian Ocean via wave processes and causes a weakening and shoaling of the AC. Our results highlight the importance of remote processes in future ocean circulation changes.

1. Introduction

The Agulhas Current (AC) is the strongest western boundary current in the Southern Hemisphere and plays a critical role in the climate system (Beal et al., 2011). Originating from the Mozambique Channel and the South-east Madagascar Current, the AC transports around 76 Sv of warm water poleward (Beal et al., 2015; McMonigal et al., 2020). After passing the southern tip of Africa, the AC retroflects back into the Indian Ocean, with a significant amount of warm and salty water leaked into the Atlantic, acting as a warm route for the surface branch of the Atlantic Meridional Overturning Circulation (AMOC; Biastoch, Lutjeharms, et al., 2008; Rühls et al., 2013; Wang et al., 2016). Thus, the AC plays a crucial role in connecting major ocean basins and could potentially modulate the variability of the global ocean overturning circulation (Biastoch, Böning, & Lutjeharms, 2008; Talley, 2013; Weijer et al., 2002). In this study, we focus on the centennial responses of the AC volume transport to the increasing greenhouse gas forcing. Climate models consistently project the AC volume transport to decline in the 21st century (Sen Gupta et al., 2021; Stellema et al., 2019). Here we argue that remote processes in the North Atlantic are as important as local processes (e.g., surface wind) in determining the AC changes in a warming climate.

The global ocean circulation is expected to change in a warming climate, but not uniformly (Chen et al., 2019; Hu et al., 2020; Peng et al., 2022). In response to the increasing greenhouse gas forcing, the atmospheric circulation evolves with a poleward shift in the westerly wind in both hemispheres (Chen et al., 2008; Goyal et al., 2021). This shift in the westerly wind leads to a poleward expansion of the subtropical gyre circulation (Yang et al., 2020) and could potentially contribute to a weakening of the western boundary currents (Ma et al., 2020). This wind changes have been suggested to contribute to the weakening of AC in a warming climate (Stellema et al., 2019).

© 2023 The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.




Figure 1. A schematic of the Indian Ocean circulation and its changes. The background color represents changes in wind stress curl from 2006 to 2100 in CESM Large Ensemble. The red line denotes the cross-section for evaluating the AC volume transport. The green arrows represent the direction of time-mean circulation and the purple arrows represent the changes. The light purple arrows around the African continent indicate the pathway by which the AMOC affect the AC.

On the other hand, as the climate warms, sea surface temperature increases and leads to a stronger upper-ocean density stratification (Capotondi et al., 2012; Li et al., 2020). The stronger density stratification limits the vertical momentum input into the subsurface ocean, causing a spin-up of the upper ocean circulation and a spin-down of the subsurface ocean circulation (Chen et al., 2019; Peng et al., 2022; Sun et al., 2013; Zhang et al., 2017). This stratification change has been invoked in explaining the strengthening of the Kuroshio Current (Chen et al., 2019).

Previous studies on the long-term weakening of the AC transport have primarily focused on changes in the local surface wind forcing and the Indonesian Throughflow (ITF) (Ma et al., 2020; Stellema et al., 2019). In the mean state, the upper-ocean volume transports across the boundaries of a basin are balanced and, thus, the AC volume transport can be expressed as a combination of the ITF volume transport and the wind-driven Sverdrup transport across the Indian Ocean (green arrows in Figure 1), that is,

$$V_{AC} = V_{ITF} + \int_{x_w}^{x_e} \frac{\text{curl } \bar{\tau}}{\beta \rho_0} dx, \quad (1)$$

where V_{AC} and V_{ITF} represent the volume transport of the AC (positive for southward transport) and ITF (positive for transport into the Indian Ocean), respectively, $\text{curl } \bar{\tau}$ is the curl of surface wind stress, β is the meridional gradient of the Coriolis parameter, ρ_0 is the reference seawater density. Critically, the interior diapycnal upwelling in the Indian Ocean might also contribute to the volume transport of AC, but it is small, ~ 1 Sv, compared to the upper-ocean transport (Text S1 in Supporting Information S1). In a warming climate, the ITF volume transport is projected to decline (Feng et al., 2017; Sen Gupta et al., 2016) and the Indian Ocean wind stress curl is expected to develop a negative anomaly at around 30°S, as the Southern Hemisphere westerly wind shifts southward, both contributing to a weaker AC (purple arrows in Figure 1). However, as we estimated in Section 2, changes in the ITF and Indian Ocean wind stress forcing explain less than 60% of the AC volume transport decrease in most of the climate models examined in this study. This discrepancy arises because the upper-ocean volume transport balance in Equation 1 is not satisfied in transient responses due to a volume transport convergence that exists in balance with the deepening of isopycnal surfaces, following a weakening of the AMOC (Sun et al., 2020, 2022).

Due to surface warming and freshening in the high-latitude North Atlantic, the AMOC is expected to slow down in the 21st century with a reduced formation of North Atlantic Deep Water (NADW; Cheng et al., 2013; Weijer et al., 2020). The reduction in NADW formation causes a deepening of isopycnals in the high-latitude North Atlantic. The isopycnal deepening signal propagates into the South Atlantic and Indian Ocean via Kelvin waves and Rossby waves (Huang et al., 2000), leading to geostrophically-balanced, volume transport anomalies that are opposite between the Atlantic and Indo-Pacific basins (Sun et al., 2020; Sun & Thompson, 2020). Building on this idea, we show that the isopycnal deepening along the western boundary of the Indian Ocean could modify the baroclinic structure of the AC and contribute substantially to the projected AC weakening in the 21st century.




Figure 2. Comparison of different processes in contributing to the AC volume transport reduction in CESM and the reduced gravity model. (a) Evolution of the ensemble- and annual-mean AC volume transport anomaly (blue solid line) and its linear trend (blue dashed line) in CESM-LENS during 2006–2100. The sum of the annual-mean Sverdrup transport anomaly and ITF volume transport anomaly (purple solid line) as well as its linear trend (purple dashed line) during 2006–2100 are plotted for comparison. (b) The decline of the AC volume transport from 2006 to 2100 in CESM and the contributions due to changes in Sverdrup transport and ITF volume transport. The ITF volume transport reduction due to Indo-Pacific wind changes is estimated based on the Island Rule constraints. (c) The AC volume transport anomaly (Sv) simulated by the reduced gravity model (blue line) as compared to the CESM-LENS member (gray line) and its linear fitting (red line). The AC volume transport anomaly forced by the AMOC alone and the Indo-Pacific wind alone is plotted as a black and yellow line. (d) The decline of the AC volume transport (Sv) simulated by the reduced gravity model from 2006 to 2100. Changes in panels (b) and (d) are calculated as the linear trend during 2006–2100, multiplied by 94 years.

Our main approach is to use a hierarchy of ocean and climate models to investigate the physical processes responsible for the projected rapid weakening of the AC in a warming climate. We first use a suite of coupled climate model simulations to estimate how much of the AC weakening in a warming climate can be explained by changes in local surface wind and ITF, as described in Section 2. We then use a reduced gravity model and an ocean-only general circulation model to illustrate the key processes for the AMOC to influence the AC volume transport in Sections 3 and 4. We summarize the main results in Section 5.

2. AC Changes in Climate Models

To evaluate how much of the AC changes in a warming climate can be explained by local vs remote processes, we use model output from the Community Earth System Model Large Ensemble (CESM-LENS) (Kay et al., 2015) under the Representative Concentration Pathway 8.5 (RCP8.5) high-emission scenario from 2006 to 2100. We calculate the ensemble mean of the 40 CESM-LENS members. The AC volume transport across 32.5°S (red line in Figure 1), integrated from the sea surface to 2,000 m depth and averaged between 2006 and 2015, is 69.2 Sv, which is close to observations (Beal et al., 2015; Bryden et al., 2005; Gunn et al., 2020; McMonigal et al., 2020). From 2006 to 2100, the AC volume transport declines substantially by about 16.5 Sv (Figure 2a), or ~24%.

Changes in the Indian Ocean surface wind and ITF cannot fully explain the projected AC transport decrease (cf., Ma et al., 2020). The negative wind stress curl anomaly in the Indian subtropics corresponds to a weakening of the interior Sverdrup transport and thus a decline of the southward volume transport by the AC [Equation 1 and Figure 1]. At 32.5°S, we estimate that this change contributes 4.4 Sv to the AC volume transport reduction from 2006 to 2100 (Figure 2b). The ITF volume transport declines by about 4.2 Sv from 2006 to 2100 (Figure 2b). Taken together, changes in the Indian Ocean surface wind and ITF volume transport explain 52% of the AC weakening (pink line vs. blue line in Figure 2a). Critically, part of the ITF volume transport reduction may be

Table 1
Description of the Reduced Gravity Model Experiments

Experiments	Indian wind	Pacific wind	AMOC forcing
Ind_wind	CESM	Control	Control
Pac_wind	Control	CESM	Control
AMOC_ForCing	Control	Control	CESM
AllForcing	CESM	CESM	CESM

Note. “CESM” means the forcing evolves as in CESM during 2006–2100; “Control” means the forcing is fixed at year 2006 as in the control run.

driven by remote changes in the North Atlantic (Sun & Thompson, 2020). We estimate the ITF volume transport reduction due to changes in local Indo-Pacific surface wind based on the Island Rule constraint and conclude that local processes only account for 1.6 Sv of the ITF reduction (Figure S1 in Supporting Information S1, Godfrey, 1989), with the remaining 2.6 Sv related to the weakening of the AMOC (Sun & Thompson, 2020). We note that the Island Rule calculation is sensitive to the choice of integral pathway, but the conclusion should be robust qualitatively.

Similar conclusions are drawn for models that participate the Coupled Model Intercomparison Project, Phase 6 (CMIP6) (Eyring et al., 2016). We examined the circulation changes in 20 CMIP6 climate models under the high-end emission scenario (“SSP585”) from 2015 to 2100. As the climate warms, all

20 models exhibit a declining trend in the AC volume transport, ranging from 10 to 35 Sv. In 12 models, changes in the Indian Ocean wind and ITF transport explain less than 60% of the AC weakening (Figure S2 in Supporting Information S1). Only in five of the 20 models, changes in the Indian Ocean wind and ITF transport explain more than 80% of the AC weakening. A brief discussion on the inter-model spread is included in Section 5. Nonetheless, these results suggest that there is another process in play that influence the transient responses of the AC volume transport. Here we argue it is the weakening of the AMOC that accounts for the discrepancies between the simulated AC volume transport changes and previous expectations based on the mean-state upper-ocean volume transport balance (Equation 1). In the next section, we illustrate the processes by which the AMOC weakening contributes to the AC volume transport decrease using a 1.5-layer reduced gravity model.

3. AC Changes in a Reduced Gravity Model

In this section, we use a 1.5-layer reduced gravity model to investigate the role of AMOC weakening and Indo-Pacific wind changes in driving the decline of the AC volume transport, projected for the 21st century. Reduced gravity models have proven in previous studies to be useful tools in guiding theoretical discussions of the physical processes that govern the large-scale circulation (Allison et al., 2011; Chen et al., 2014; Johnson & Marshall, 2004; Sun et al., 2020).

We use the 1.5-layer reduced gravity model developed in Sun and Thompson (2020). The model is an idealized representation of the wind-driven gyre circulation and the upper branch of the global ocean overturning circulation. The model domain includes three idealized ocean basins representative of the Atlantic, Indian, and Pacific, connected by a zonally re-entrant channel that represents the Southern Ocean (Figure S3a in Supporting Information S1). The model evolves the upper layer thickness h in the ocean. Surface water mass transformation in the Southern Ocean is represented as a relaxation of the upper layer thickness to 10 m close to the southern boundary. The relaxation timescale increases from 10 days at the southern boundary to 100 days at 62°S. As such, the Southern Ocean overturning circulation is allowed to evolve as the global-mean interface depth changes in the northern basins. Interior diapycnal mixing is parameterized as a diapycnal upwelling velocity, $w_{\text{diap}} = \kappa/h$, with $\kappa = 2.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$. The interior diapycnal mixing contributes about 1 Sv of diapycnal upwelling, close to that in CESM-LENS (Text S1 in Supporting Information S1). The unresolved mesoscale eddies are parameterized as a thickness diffusion with the diffusivity $K_{\text{GM}} = 1,000 \text{ m}^2 \text{ s}^{-1}$ (Gent & McWilliams, 1990). We force the reduced gravity model with annual-mean surface wind stress and NADW formation rate derived from the first member of CESM-LENS. The NADW formation rate in CESM is calculated as the maximum value of the AMOC at 40°N and is prescribed in the reduced gravity model as a downward velocity across the layer interface close to the northern boundary of the Atlantic basin. A detailed description of the model configuration is provided in Text S2 in Supporting Information S1. As a control, we integrate the reduced gravity model, forced by the annual-mean surface wind stress (Figure S3a in Supporting Information S1) and NADW formation rate from CESM at year 2006, for 4,000 years to reach an approximately steady state. Using control as initial conditions, we conduct another four simulations, summarized in Table 1, to quantify the contributions to the weakening of the AC due to changes in the Indo-Pacific wind and AMOC (Figure S4 in Supporting Information S1).

When forced with both the Indo-Pacific wind and NADW formation rate that evolve as in CESM during 2006–2100 (“ALLFORCING”), the reduced gravity model approximately reproduces the AC volume transport changes (compare the blue and red lines in Figure 2c). This suggests that the reduced gravity model captures the key




Figure 3. Connecting the shoaling of the AC to AMOC weakening. (a) Changes in the AMOC streamfunction from 2006 to 2100 in the CESM-LENS (background color), calculated as 94 times the linear trend. The black contours denote the time-mean AMOC streamfunction averaged between 2006 and 2100 (positive for clockwise overturning). (b) Changes in in-situ density (kg/m^3) along the AC cross-section (red line in Figure 1) from 2006 to 2100 in CESM-LENS. The density anomaly on the right boundary ($x = 200 \text{ km}$) is subtracted to highlight the offshore density gradient. The gray patch in the left side denotes the coast of Africa. (c) Velocity changes (m/s) in the AC section from 2006 to 2100 in CESM-LENS (negative for AC slowdown). The black contours denote the time-mean velocity of AC averaged between 2006 and 2100. The gray patch in the left side denotes the coast of Africa. (d–f) Same as above but for the “AMOC weakening” experiment carried out with MITgcm. The anomalies are calculated as the difference between the “AMOC weakening” run and control run, averaged over the last 20 years.

dynamics in driving the projected AC weakening in CESM. When we only evolve the Indian Ocean wind as in CESM and fix the Pacific wind and the AMOC as in Control (“IND_WIND”), the reduced gravity model only simulates a weakening of the AC by 4.4 Sv (Figure 2d), consistent with the estimated Sverdrup transport changes in CESM (Figure 2b). When we only evolve the Pacific surface wind as in CESM during 2006–2100 (“PAC_WIND”), the reduced gravity model simulates a weakening of the AC by 1.1 Sv. This number is much smaller than the decline of ITF volume transport ($\sim 4.2 \text{ Sv}$) in CESM, but it is close to our estimate from changes in the Indo-Pacific wind based on the Island Rule constraints (1.6 Sv in Figure 2b). The discrepancy arises because much of the projected ITF changes in CESM are driven by the weakening of the AMOC (Sun & Thompson, 2020), which is not represented in “PAC_WIND.” To identify the impacts from the weakening of the AMOC, we carry out “AMOC_FORCING,” in which we fix the surface wind stress and only evolve the NADW formation rate as in CESM. The weakening of the AMOC leads to a decline of the AC volume transport by about 11 Sv (Figures 2c and 2d). Taken together, changes in the Indo-Pacific wind and AMOC explains the decline of the AC volume transport. The impact of AMOC exceeds the contributions due to surface wind changes, highlighting the role of remote processes in the projected decline of the AC volume transport.

The remote North Atlantic processes influence the AC volume transport via wave propagations. As the AMOC slows down in a warming climate, isopycnals deepen in the high-latitude North Atlantic (Sun et al., 2020). The deepening signal propagates as Kelvin waves along the coast and the equator to the western boundary of the Indian Ocean (Figure S5 in Supporting Information S1; Huang et al., 2000). The isopycnal deepening along the Indian Ocean western boundary corresponds to a northward geostrophic transport anomaly, that is, a slowdown of the AC (Figure S6 in Supporting Information S1 for the reduced gravity model).

Despite the idealized nature of the reduced gravity model, the physical processes linking changes in the AMOC to the AC discussed above appears to be also relevant in more realistic simulations (Figures 2 and 3). In CESM, there is a positive anomaly in the offshore density gradient across the AC, most pronounced at 1–2 km depths, consistent with a deepening of isopycnals along the western boundary (Figure 3b and Figure S7 in Supporting

Information S1). This depth range aligns well with the structure of the AMOC weakening in CESM, which also peaks at 1–2 km (Figure 3a). The density anomaly corresponds to a slow-down of the AC through geostrophy.

4. Shoaling of the AC in a Warming Climate

The weakening of the AC is accompanied by a change in its depth structure. In CESM-LENS, the velocity field of the AC throughout the upper 2 km exhibits a decreasing trend, which peaks at about 1 km depth (Figure 3c) and represents a shoaling of the AC depth. Defined as the maximum depth of the 1 cm/s velocity contour, the depth of AC is shoaled from 1,700 m at 2006 to 1,300 m at 2100 (Figure S8 in Supporting Information S1). Similar shoaling of the AC is present in other climate models that participate CMIP6 (Figures S9 and S10 in Supporting Information S1). Both the AMOC weakening and the stronger surface stratification in a warming climate may contribute to this shoaling of the AC (e.g., Peng et al., 2022). Here we use a suite of ocean-only simulations to show that this shoaling of the AC is mainly driven by the remote AMOC changes.

We use the Massachusetts Institute of Technology General Circulation Model (MITgcm; Marshall et al., 1997) to carry out the ocean-only simulations. The model uses the same geometry as the reduced gravity model. The model domain has 30 vertical layers ranging from 20 m at the top to 250 m at the bottom. There is only one thermodynamic variable, temperature, for simplicity. We perturb the AMOC through a warming in the high-latitude North Atlantic. A detailed discussion of the model setup is provided in Text S3 in Supporting Information S1. As a control run, we run the model for 2,000 years to reach an approximately steady state. Using the control run as initial conditions, we evaluate the role of remote and local processes in driving the AC changes. Two experiments are designed: “AMOC weakening” and “Indian Ocean warming.”

In “AMOC weakening,” we apply a 3°C sea surface warming in the high-latitude North Atlantic from 67.5°N to 71.5°N. This surface warming strengthens the upper-ocean stratifications in the North Atlantic and leads to a decline of the AMOC strength by 6 Sv (Figure 3d and Figure S12 in Supporting Information S1). Similar to CESM, there is also a positive anomaly in the offshore density gradient across the AC. This density gradient anomaly is most pronounced at 1–2 km depths and corresponds to a deepening of the isopycnal driven by an AMOC weakening (Figures 3d and 3e). The AMOC weakening causes the AC volume transport to decrease by a similar magnitude of around 6 Sv (Figure S12 in Supporting Information S1), largely consistent with the reduced gravity model in Section 3. The velocity field shows a decreasing trend through the upper 1.6 km. The decreasing trend peaks at around 0.8 km depth, slightly shallower than CESM, which is not unexpected considering their different model configurations. The depth of AC, defined as the maximum depth of 1 cm/s velocity contour, shoals by about 200 m in the “AMOC weakening” (Figure S11 in Supporting Information S1).

The surface warming in the Indian Ocean could also contribute to the subsurface weakening of the AC (Peng et al., 2022). Here we test this idea by applying a 3°C warming in the Indian Ocean (“Indian Ocean warming” run). Consistent with previous studies, the surface warming leads to a spin-up of the upper ocean circulation and a spin-down of the subsurface gyre circulation (Figure S13 in Supporting Information S1). However, these changes are confined to the upper 600 m and does not appear to affect the AC depth by our definition (Figure S11 in Supporting Information S1). When integrated vertically, the surface spin-up and subsurface spin-down also cancels out, leading to minimal changes in the AC volume transport (Figure S12 in Supporting Information S1).

We also examined the influence of the Southern Hemisphere (SH) westerly wind in the AC changes. The SH westerly wind is projected to strengthen in a warming climate (Goyal et al., 2021) and may also contribute to the AC changes. However, our results suggest that a strengthening of the SH westerly wind by 10%, comparable to the projected changes in CMIP6, actually strengthens the AC from 300 to 800 m depth (Figure S14 in Supporting Information S1).

5. Summary and Discussions

Climate models consistently project a weakening of the AC during the 21st century. Previous studies typically attributed this weakening to changes in the local surface wind stress or/and ITF volume transport. However, our analysis suggests these changes cannot fully explain the AC changes in climate model simulations. Here we revisit the physical processes in driving the AC weakening and highlight the role of AMOC weakening as an additional process in contributing to the AC changes. In response to the increasing greenhouse gas forcing, the

AMOC slows down and modifies the baroclinic structure of the AC, communicated between basins by waves. This process leads to a weakening and shoaling of AC in a warming climate during the 21st century.

The 21st century projections for the AC transport stand in contrast with the historical changes. During the historical period, it remains debatable whether the AC strength has changed significantly in response to greenhouse gas forcing (cf., Beal & Elipot, 2016). Recently, using a hierarchy of ocean and climate models, Zhang et al. (2023) showed that the AC is characterized by substantial decadal and multi-decadal variabilities, which complicated the interpretation of the AC responses based on a short observational period during the historical period.

Although the dynamical connection between the AMOC and the AC is robust, our results suggest significant inter-model spread in how much the AMOC explains the projected AC changes in the 21st century (e.g., Figure S2 in Supporting Information S1). This may arise from the inter-model spread in the projected AMOC weakening during the 21st century (e.g., Bellomo et al., 2021; Bonan et al., 2022) or the inter-model spread in the projected wind changes (e.g., Sen Gupta et al., 2016).

We emphasize that the remote impact of AMOC weakening on the AC is a transient process that is mostly highlighted on decadal to centennial timescales. On longer timescales, as the isopycnal deepens in the Indo-Pacific basin, the Southern Ocean overturning circulation will gradually play a bigger role in balancing the AMOC changes on multi-centennial timescales (Sun et al., 2020). Eventually, the AC will be in balance with the local Sverdrup transport and ITF volume transport again (Equation 1) in steady state.

The rapid weakening of AC might play an important role in the climate system. As one of the strongest western boundary currents, the AC transport a large amount of heat poleward. In a warming climate, the weakening of AC could induce a heat convergence and contribute to the warming of the Indian Ocean (McMonigal et al., 2022; Sun et al., 2022). Additionally, changes in the AC system in a warming climate might play an important role in modulating the strength of AMOC by changing the salt budget in the Atlantic basin (Beal et al., 2011; Biastoch et al., 2009; Weijer et al., 2002). However, due to the coarse resolution in most of the climate models, the simulated Agulhas Leakage is likely unrealistic, which might affect our understanding of how the Agulhas Leakage changes, following a weakening of the AC, may feed back to the overturning circulation (Weijer and van Sebille, 2014). In the future, more state-of-the-art high-resolution climate models are needed to evaluate the potential feedbacks to the Atlantic overturning due to changes in the AC system.

Data Availability Statement

The CESM large ensemble data were downloaded from Climate Data Gateway at NCAR (<https://www.earthsystemgrid.org/>). The CMIP6 data were downloaded from the Earth System Grid Federation node (<https://esgf-node.llnl.gov/search/cmip6/>). The reduced gravity model is available at the online open access repository figshare (<https://doi.org/10.6084/m9.figshare.12903086>) under a “CC BY 4.0” licence. The MITgcm experiments source code and the data used to generate figures are now available in figshare (<https://doi.org/10.6084/m9.figshare.21802770>).

References

- Allison, L. C., Johnson, H. L., & Marshall, D. P. (2011). Spin-up and adjustment of the Antarctic Circumpolar Current and global pycnocline. *Journal of Marine Research*, 69(2), 167–189. <https://doi.org/10.1357/002224011798765330>
- Beal, L. M., De Ruijter, W. P., Biastoch, A., Zahn, R., Group, S. W. I. W., Hermes, J., et al. (2011). On the role of the Agulhas system in ocean circulation and climate. *Nature*, 472(7344), 429–436. <https://doi.org/10.1038/nature09983>
- Beal, L. M., & Elipot, S. (2016). Broadening not strengthening of the Agulhas Current since the early 1990s. *Nature*, 540(7634), 570–573. <https://doi.org/10.1038/nature19853>
- Beal, L. M., Elipot, S., Houk, A., & Leber, G. M. (2015). Capturing the transport variability of a western boundary jet: Results from the Agulhas current time-series experiment (ACT). *Journal of Physical Oceanography*, 45(5), 1302–1324. <https://doi.org/10.1175/jpo-d-14-0119.1>
- Bellomo, K., Angeloni, M., Corti, S., & von Hardenberg, J. (2021). Future climate change shaped by inter-model differences in Atlantic meridional overturning circulation response. *Nature Communications*, 12(1), 3659. <https://doi.org/10.1038/s41467-021-24015-w>
- Biastoch, A., Böning, C. W., & Lutjeharms, J. (2008). Agulhas leakage dynamics affects decadal variability in Atlantic overturning circulation. *Nature*, 456(7221), 489–492. <https://doi.org/10.1038/nature07426>
- Biastoch, A., Böning, C. W., Schwarzkopf, F. U., & Lutjeharms, J. R. (2009). Increase in Agulhas leakage due to poleward shift of Southern Hemisphere westerlies. *Nature*, 462(7272), 495–498. <https://doi.org/10.1038/nature08519>
- Biastoch, A., Lutjeharms, J. R. E., Böning, C. W., & Scheinert, M. (2008). Mesoscale perturbations control inter-ocean exchange south of Africa. *Geophysical Research Letters*, 35(20), L20602. <https://doi.org/10.1029/2008gl035132>
- Bonan, D. B., Thompson, A. F., Newsom, E. R., Sun, S., & Rugenstein, M. (2022). Transient and equilibrium responses of the Atlantic overturning circulation to warming in coupled climate models: The role of temperature and salinity. *Journal of Climate*, 35(15), 1–42. <https://doi.org/10.1175/jcli-d-21-0912.1>

Acknowledgments

This research is supported by the National Natural Science Foundation of China (42225601, 42076009) and Fundamental Research Funds for the Central Universities (202072001). S. Sun is supported by Florida State University. Z. Chen is partly supported by the Taishan Scholar Funds (tsqn201812022). The authors thank all anonymous reviewers for providing constructive comments.

- Bryden, H. L., Beal, L. M., & Duncan, L. M. (2005). Structure and transport of the Agulhas current and its temporal variability. *Journal of Oceanography*, 61(3), 479–492. <https://doi.org/10.1007/s10872-005-0057-8>
- Capotondi, A., Alexander, M. A., Bond, N. A., Curchitser, E. N., & Scott, J. D. (2012). Enhanced upper ocean stratification with climate change in the CMIP3 models. *Journal of Geophysical Research*, 117(C4), C04031. <https://doi.org/10.1029/2011jc007409>
- Chen, C., Wang, G., Xie, S.-P., & Liu, W. (2019). Why does global warming weaken the Gulf Stream but intensify the Kuroshio? *Journal of Climate*, 32(21), 7437–7451. <https://doi.org/10.1175/jcli-d-18-0895.1>
- Chen, G., Lu, J., & Frierson, D. M. W. (2008). Phase speed spectra and the latitude of surface westerlies: Interannual variability and global warming trend. *Journal of Climate*, 21(22), 5942–5959. <https://doi.org/10.1175/2008jcli2306.1>
- Chen, Z., Wu, L., Qiu, B., Sun, S., & Jia, F. (2014). Seasonal variation of the south equatorial current bifurcation off Madagascar. *Journal of Physical Oceanography*, 44(2), 618–631. <https://doi.org/10.1175/jpo-d-13-0147.1>
- Cheng, W., Chiang, J. C. H., & Zhang, D. (2013). Atlantic meridional overturning circulation (AMOC) in CMIP5 models: RCP and historical simulations. *Journal of Climate*, 26(18), 7187–7197. <https://doi.org/10.1175/jcli-d-12-00496.1>
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K. E. (2016). Overview of the coupled model inter-comparison project phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*, 9(5), 1937–1958. <https://doi.org/10.5194/gmd-9-1937-2016>
- Feng, M., Zhang, X., Sloyan, B., & Chamberlain, M. (2017). Contribution of the deep ocean to the centennial changes of the Indonesian Throughflow. *Geophysical Research Letters*, 44(6), 2859–2867. <https://doi.org/10.1002/2017gl072577>
- Gent, P. R., & McWilliams, J. C. (1990). Isopycnal mixing in ocean circulation models. *Journal of Physical Oceanography*, 20(1), 150–155. [https://doi.org/10.1175/1520-0485\(1990\)020<0150:imicm>2.0.co;2](https://doi.org/10.1175/1520-0485(1990)020<0150:imicm>2.0.co;2)
- Godfrey, J. S. (1989). A Sverdrup model of the depth-integrated flow for the world ocean allowing for island circulations. *Geophysical & Astrophysical Fluid Dynamics*, 45(1–2), 89–112. <https://doi.org/10.1080/03091928908208894>
- Goyal, R., Sen Gupta, A., Jucker, M., & England, M. H. (2021). Historical and projected changes in the southern hemisphere surface westerlies. *Geophysical Research Letters*, 48(4), e2020GL090849. <https://doi.org/10.1029/2020gl090849>
- Gunn, K. L., Beal, L. M., Elipot, S., McMonigal, K., & Houk, A. (2020). Mixing of subtropical, central, and intermediate waters driven by shifting and pulsing of the Agulhas current. *Journal of Physical Oceanography*, 50(12), 3545–3560. <https://doi.org/10.1175/jpo-d-20-0093.1>
- Hu, S., Sprintall, J., Guan, C., McPhaden, M. J., Wang, F., Hu, D., & Cai, W. (2020). Deep-reaching acceleration of global mean ocean circulation over the past two decades. *Science Advances*, 6, eaax7727. <https://doi.org/10.1126/sciadv.aax7727>
- Huang, R. X., Cane, M. A., Naik, N., & Goodman, P. (2000). Global adjustment of the thermocline in response to deepwater formation. *Geophysical Research Letters*, 27(6), 759–762. <https://doi.org/10.1029/1999gl002365>
- Johnson, H. L., & Marshall, D. P. (2004). Global teleconnections of meridional overturning circulation anomalies. *Journal of Physical Oceanography*, 34(7), 1702–1722. [https://doi.org/10.1175/1520-0485\(2004\)034<1702:gtomoc>2.0.co;2](https://doi.org/10.1175/1520-0485(2004)034<1702:gtomoc>2.0.co;2)
- Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., et al. (2015). The community Earth system model (CESM) large ensemble project: A community resource for studying climate change in the presence of internal climate variability. *Bulletin of the American Meteorological Society*, 96(8), 1333–1349. <https://doi.org/10.1175/bams-d-13-00255.1>
- Li, G., Cheng, L., Zhu, J., Trenberth, K. E., Mann, M. E., & Abraham, J. P. (2020). Increasing ocean stratification over the past half-century. *Nature Climate Change*, 10(12), 1116–1123. <https://doi.org/10.1038/s41558-020-00918-2>
- Ma, J., Feng, M., Lan, J., & Hu, D. (2020). Projected future changes of meridional heat transport and heat balance of the Indian Ocean. *Geophysical Research Letters*, 47(4), e2019GL086803. <https://doi.org/10.1029/2019gl086803>
- Marshall, J., Adcroft, A., Hill, C., Perelman, L., & Heisey, C. (1997). A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers. *Journal of Geophysical Research*, 102(C3), 5753–5766. <https://doi.org/10.1029/96jc02775>
- McMonigal, K., Beal, L. M., Elipot, S., Gunn, K. L., Hermes, J., Morris, T., & Houk, A. (2020). The impact of meanders, deepening and broadening, and seasonality on Agulhas current temperature variability. *Journal of Physical Oceanography*, 50(12), 3529–3544. <https://doi.org/10.1175/jpo-d-20-0018.1>
- McMonigal, K., Gunn, K. L., Beal, L. M., Elipot, S., & Willis, J. K. (2022). Reduction in meridional heat export contributes to recent Indian Ocean warming. *Journal of Physical Oceanography*, 52(3), 329–345. <https://doi.org/10.1175/jpo-d-21-0085.1>
- Peng, Q., Xie, S. P., Wang, D., Huang, R. X., Chen, G., Shu, Y., et al. (2022). Surface warming-induced global acceleration of upper ocean currents. *Science Advances*, 8(16), eabj8394. <https://doi.org/10.1126/sciadv.abj8394>
- Rühs, S., Durgadoo, J. V., Behrens, E., & Biastoch, A. (2013). Advective timescales and pathways of Agulhas leakage. *Geophysical Research Letters*, 40(15), 3997–4000. <https://doi.org/10.1002/grl.50782>
- Sen Gupta, A., McGregor, S., van Sebille, E., Ganachaud, A., Brown, J. N., & Santoso, A. (2016). Future changes to the Indonesian Throughflow and Pacific circulation: The differing role of wind and deep circulation changes. *Geophysical Research Letters*, 43(4), 1669–1678. <https://doi.org/10.1002/2016gl067757>
- Sen Gupta, A., Stellega, A., Pontes, G. M., Taschetto, A. S., Vergés, A., & Rossi, V. (2021). Future changes to the upper ocean Western Boundary Currents across two generations of climate models. *Scientific Reports*, 11(1), 9538. <https://doi.org/10.1038/s41598-021-88934-w>
- Stellega, A., Sen Gupta, A., & Taschetto, A. S. (2019). Projected slow down of South Indian Ocean circulation. *Scientific Reports*, 9(1), 17705. <https://doi.org/10.1038/s41598-019-54092-3>
- Sun, S., & Thompson, A. F. (2020). Centennial changes in the Indonesian throughflow connected to the Atlantic meridional overturning circulation: The Ocean's transient conveyor belt. *Geophysical Research Letters*, 47(21), e2020GL090615. <https://doi.org/10.1029/2020gl090615>
- Sun, S., Thompson, A. F., & Eisenman, I. (2020). Transient overturning compensation between Atlantic and Indo-Pacific basins. *Journal of Physical Oceanography*, 50(8), 2151–2172. <https://doi.org/10.1175/jpo-d-20-0060.1>
- Sun, S., Thompson, A. F., Xie, S.-P., & Long, S.-M. (2022). Indo-Pacific warming induced by a weakening of the Atlantic meridional overturning circulation. *Journal of Climate*, 35(2), 815–832. <https://doi.org/10.1175/jcli-d-21-0346.1>
- Sun, S., Wu, L., & Qiu, B. (2013). Response of the inertial recirculation to intensified stratification in a two-layer quasigeostrophic ocean circulation model. *Journal of Physical Oceanography*, 43(7), 1254–1269. <https://doi.org/10.1175/jpo-d-12-0111.1>
- Talley, D. L. (2013). Closure of the global overturning circulation through the Indian, Pacific, and Southern Oceans: Schematics and transports. *Oceanography*, 26(1), 80–97. <https://doi.org/10.5670/oceanog.2013.07>
- Wang, Y., Beron-Vera, F. J., & Olascoaga, M. J. (2016). The life cycle of a coherent Lagrangian Agulhas ring. *Journal of Geophysical Research: Oceans*, 121(6), 3944–3954. <https://doi.org/10.1002/2015jc011620>
- Weijer, W., Cheng, W., Garuba, O. A., Hu, A., & Nadiga, B. T. (2020). CMIP6 Models predict significant 21st century decline of the Atlantic meridional overturning circulation. *Geophysical Research Letters*, 47(12), e2019GL086075. <https://doi.org/10.1029/2019gl086075>
- Weijer, W., De Ruijter, W. P. M., Sterl, A., & Drijfhout, S. S. (2002). Response of the Atlantic overturning circulation to South Atlantic sources of buoyancy. *Global and Planetary Change*, 34(3–4), 293–311. [https://doi.org/10.1016/s0921-8181\(02\)00121-2](https://doi.org/10.1016/s0921-8181(02)00121-2)

- Weijer, W., & van Sebille, E. (2014). Impact of Agulhas leakage on the Atlantic overturning circulation in the CCSM4. *Journal of Climate*, 27(1), 101–110. <https://doi.org/10.1175/jcli-d-12-00714.1>
- Yang, H., Lohmann, G., Krebs-Kanzow, U., Ionita, M., Shi, X., Sidorenko, D., et al. (2020). Poleward shift of the major ocean gyres detected in a warming climate. *Geophysical Research Letters*, 47(5), e2019GL085868. <https://doi.org/10.1029/2019gl085868>
- Zhang, R., Sun, S., Chen, Z., Yang, H., & Wu, L. (2023). On the decadal and multi-decadal variability of the Agulhas Current. *Journal of Physical Oceanography*. <https://doi.org/10.1175/JPO-D-22-0123.1>
- Zhang, X., Wang, Q., & Mu, M. (2017). The impact of global warming on Kuroshio Extension and its southern recirculation using CMIP5 experiments with a high-resolution climate model MIROC4h. *Theoretical and Applied Climatology*, 127(3–4), 815–827. <https://doi.org/10.1007/s00704-015-1672-y>

References From the Supporting Information

- de Lavergne, C., Madec, G., Roquet, F., Holmes, R. M., & McDougall, T. J. (2017). Abyssal ocean overturning shaped by seafloor distribution. *Nature*, 551(7679), 181–186. <https://doi.org/10.1038/nature24472>
- Ferrari, R., Mashayek, A., McDougall, T. J., Nikurashin, M., & Campin, J.-M. (2016). Turning ocean mixing upside down. *Journal of Physical Oceanography*, 46(7), 2239–2261. <https://doi.org/10.1175/jpo-d-15-0244.1>
- Ledwell, J., Watson, A., & Law, C. (1993). Evidence for slow mixing across the pycnocline from an open-ocean tracer-release experiment. *Nature*, 364(6439), 701–703. <https://doi.org/10.1038/364701a0>
- Thompson, A. F., Stewart, A. L., & Bischoff, T. (2016). A multibasin residual-mean model for the global overturning circulation. *Journal of Physical Oceanography*, 46(9), 2583–2604. <https://doi.org/10.1175/jpo-d-15-0204.1>
- Waterhouse, A., MacKinnon, J. A., Nash, J. D., Alford, M. H., Kunze, E., Simmons, H. L., et al. (2014). Global patterns of diapycnal mixing from measurements of the turbulent dissipation rate. *Journal of Physical Oceanography*, 44(7), 1854–1872. <https://doi.org/10.1175/jpo-d-13-0104.1>